

PRIMER NOTE

Improved noninvasive genotyping method: application to brown bear (*Ursus arctos*) faeces

E. BELLEMAIN*† and P. TABERLET*

*Laboratoire d'Ecologie Alpine (LECA), Génomique des Populations et Biodiversité, CNRS UMR 5553, Université Joseph Fourier, BP 53, F-38041 Grenoble Cedex 9, France, †Department of Ecology and Natural Resource Management, Agricultural University of Norway, Postbox 5014, N-1432 Ås, Norwegian Institute for Nature Research, Tungasletta 2, N-7485 Trondheim, Norway

Abstract

We redesigned new microsatellite primers and one sex-specific primer for amplification of faecal DNA from brown bears (*Ursus arctos*). We also combined a semi-nested polymerase chain reaction (PCR) with a newly developed multiplex preamplification method in order to increase the quality of the amplified DNA fragments. In comparison with a conventional PCR approach, the genotyping error rate was substantially reduced and the amplification rate was increased. This new approach could be transposed to other species where conventional PCR methods experience low success due to limited DNA concentration and/or quality.

Keywords: microsatellites, multiplex preamplification method, noninvasive samples, sex-specific primer, *Ursus arctos*

Received 24 March 2004; revision accepted 4 May 2004

Noninvasive genetic techniques have received much interest in the last few years. The use of DNA from faeces, hairs, urine or feathers as an individual genetic tag can provide useful information for population monitoring and the estimation of important genetic parameters. However, the amplification of DNA extracted from noninvasive samples is often problematic because the DNA is usually degraded and present in low quantity. This can lead to scoring errors such as allelic dropout or false alleles and will often produce incorrect genotypes (Taberlet *et al.* 1996; Taberlet & Luikart 1999; Smith *et al.* 2000). These problems are now well understood, and different methods have been proposed to limit genotyping errors and their impacts on the subsequent analyses (Schwartz *et al.* 1999; Mills *et al.* 2000; Miller *et al.* 2002; Paetkau 2003; Piggott *et al.* 2004).

Here, we combined a two-step PCR approach, or 'semi-nested PCR' with the newly defined 'multiplex preamplification method' (Piggott *et al.* 2004) to amplify DNA from brown bear (*Ursus arctos*) faecal samples, with the aim to increase the quality of the multilocus genotypes.

For all collected bear faeces, DNA was extracted using the Qiamp DNA Stool® kit (Qiagen), especially designed for this type of material. All extractions were carried out in

a room dedicated only to processing hairs and faeces. The final volume of the DNA extract was 150 µL. Negative controls were included in all extractions batches in order to check for exogenous DNA contamination. Each DNA extract was first screened with one microsatellite (G10P; Paetkau & Strobeck 1994) acting as a species-diagnostic marker. PCR products were separated on an agarose gel and viewed, using ethidium bromide, under UV light. Only the DNA samples that showed a fluorescent DNA band were kept for subsequent analyses.

Based on genetic data from more than 700 individuals typed for 18 or 19 microsatellite loci (Waits *et al.* 2000; E. Bellemain, unpublished data), we selected the six most informative for identifying individuals, i.e. those with the lowest probability of identity between unrelated individuals (PI; Paetkau & Strobeck 1994) and between siblings (PIsibs; Waits *et al.* 2000) (Table 1). Most of the original primers were redesigned in order to obtain smaller amplified fragments and more similar annealing temperatures. We also redesigned new sex-identification primers (Table 1), amplifying an 80 bp fragment more specific to carnivores (i.e. avoiding amplifications from most of the prey DNA co-extracted from a bear faeces).

We combined the multiplex preamplification method (Piggott *et al.* 2004) with a semi-nested PCR (Fig. 1; see Ulrich *et al.* 1993). The first step involved a multiplex

Correspondence: E. Bellemain. Fax: +33 (0)476514279; E-mail: eva.bellemain@ujf-grenoble.fr

Table 1 External and internal primers for *Ursus arctos* used for the two steps of the multiplex preamplification method

Locus	Microsatellite sequencet	Fluorescence (internal primer)	Primer's final concentration (µM)	Allele size range (bp)	PI (Paetkau & Strobeck 1994)	Plsibs (Waits <i>et al.</i> 2001)	GenBank Accession no.	Redesigned from or reference
Mu10	F*: ATTCAGATTTCATCACTTGACA R*: CCATAAATGGTAGCATGAGC	FAM	0.5	113–129	6.757e-02	3.652e-01	Y09642	Taberlet <i>et al.</i> 1997
Mu23	RI*: TCAGCATAGTTACACAATCTCC F: GCCTGTGTCTATTATCC R: TTGCTTGCCCTAGACCACC	NED	0.6	141–153	1.247e-01	4.364e-01	Y09645	Taberlet <i>et al.</i> 1997
Mu50	RI*: TAGACCACCACCAAGGCATCAG F*: GTCTCTGTCAATTCGCCATC R*: GAGCAGGAACAATGTAAGATG	FAM	0.4	92–102	1.313e-01	4.292e-01	Y09647	Taberlet <i>et al.</i> 1997
Mu51	RI*: AACCTGGAACAAAAATTAACAC F*: AGCCAGATCCTAAGAGACCT R: GAAAGGTTAGATGGAAGAGATG	HEX	0.6	105–125	8.138e-02	3.787e-01	Y09648	Taberlet <i>et al.</i> 1997
Mu59	RI*: AAAGAGAAGGACAGAGGTA F: GCTCCTTTGGACATGTAA R: AGTGTGTTGTTGGTCTGTG	NED	0.5	96–120	1.220e-01	4.183e-01	Y09649	Taberlet <i>et al.</i> 1997
G10L	RI*: TGACTGTCACCAGCAGGAG F*: CAGACAGGATATTGACATGTA R*: GATACAGAAACCTACCCTGCC	HEX	0.5	138–156	1.204e-01	4.178e-01	U22088	Paetkau & Strobeck 1994
SRY	FI: ACTGATTTTATTCACATTTCCC F: GAACGCATCTTGGTGTGGTC R: CGGTATTTATAGTCGGGTAGTT RI: TGATCTCTGAGTTTTCATTTG	HEX	0.5	75 for males only	—	—	X74007	Taberlet <i>et al.</i> 1997

+F and R represent forward and reverse external primers. FI and RI represent forward and reverse internal primers. *Indicates the microsatellite sequences that were redefined.

PI: probability of identity between unrelated individuals.

Plsibs: probability of identity between siblings.




Fig. 1 General principle of the multiplex preamplification method combined with a semi-nested PCR (the figure shows three replicates out of eight).

preamplification that simultaneously amplified all loci with external primers (Fig. 1). Each amplification was repeated eight times (multi-tubes approach; Taberlet *et al.* 1996). Aliquots from the preamplification were then used as templates in another round of PCRs. In this second step, we simultaneously amplified two or three loci (Mu10/G10L; Mu23/Mu50; SRY/Mu51/Mu59) using the internal fluorescent primers together with the appropriate external primers. The fluorescent PCR products can be electrophoresed together on the single lane.

PCR preamplifications were prepared in a 25 μL volume containing 5 μL template DNA, 0.1 mM of each dNTP, 0.01 μM of each primer, 2 mM MgCl_2 , 0.5 U AmpliTaq Gold Polymerase (Applied Biosystems) and 1 \times Taq buffer (containing 100 mM Tris-HCl, pH 8.3, 500 mM KCl, according to the manufacturer's specifications; Applied Biosystems). For the second step, PCR amplifications were prepared in a 13 μL volume containing 3 μL preamplified product, 0.1 mM of each dNTP, 0.4–0.6 μM of each primer (depending on the primer, see Table 1), 2 mM MgCl_2 , 0.5 U AmpliTaq Gold Polymerase (Applied Biosystems) and 1 \times Taq buffer (as described above). Amplifications were performed in a GeneAmp PCR system 9600 (Applied Biosystems) with the following conditions: 10 min at 95 $^\circ\text{C}$, 25 cycles for the preamplification step and 35 cycles for the second amplification step, composed of 30 s denaturing at 95 $^\circ\text{C}$, 30 s annealing

at 60 $^\circ\text{C}$, 1-min extension at 72 $^\circ\text{C}$, and as a final extension step, 7 min at 72 $^\circ\text{C}$. The PCR products from the second step were mixed (3 μL of Mu10/G10L; 3 μL of Mu23/Mu50 and 4 μL of SRY/Mu51/Mu59). One microlitre of this multiplex was added to a 10 μL mix of formamide and ROX 350 (10:0.2), and then loaded on an automatic sequencer ABI3100 (Applied Biosystems). Electrophoresis was run for 3 h with a POP4 polymer. Microsatellite patterns were examined with GENEMAPPER version 3.0 (Applied Biosystems).

The PI for the six microsatellites was low: $\text{PI} = 1.38 \times 10^{-6}$; $\text{PIsibs} = 4.52 \times 10^{-3}$, which allowed a reliable individual identification. We compared the results obtained from this multiplex approach with a conventional approach (same amplification conditions and primers as described for the second step of the multiplex approach, with 55 cycles) for eight samples and eight replicates for each locus. The amplification rate was improved substantially (Wilcoxon ranked test; $P = 0.058$) and the genotyping error (false alleles or allelic dropouts) rate was much reduced (Wilcoxon ranked test; $P = 0.063$). In addition, the DNA typing profiles were much clarified using the new multiplex method (data not shown). Next, we diluted the DNA extract from four to 12 times to simulate more limiting conditions. The genotyping error rate was from two to 10 times lower with the multiplex preamplification method than with the conventional method.

We applied this multiplex approach to more than 1900 faecal samples collected in central Sweden for estimating brown bears population size (Bellemain *et al.* 2004). A preliminary study showed that the reliability of the genotyping was not significantly different when using four or eight replicates, probably because template DNA was not in too limiting concentration in the extracts. This might be due to good field climatic conditions leading to good DNA preservation in the field before the sampling. As a consequence, we used only four replicates for this study and were able to successfully genotype 75% of the samples. To ensure the reliability of the results, we randomly repeated 5% of the genotyping and obtained an error rate of less than 2%, including the sex identification (Bellemain *et al.* 2004).

The newly defined primers, together with the PCR strategy described here, have the potential to substantially improve genotyping success rate and reduce genotyping errors. Moreover, the strategy of combining the multiplex preamplification method and a semi-nested PCR could be transposed to other species where conventional PCR approaches yield low success due to limiting DNA concentration and/or quality.

Acknowledgements

This work was financed by the Norwegian Directorate for Nature Management, the Swedish Association for Hunting and Wildlife Management, the Swedish Environmental Agency, WWF Sweden and the Orsa Communal Forest. We thank Carole Poillot, Ludovic Gielly, Delphine Rioux and Christian Miquel for help in the laboratory work. We are grateful to Sven Brunberg from the Scandinavian Brown Bear Research Project for coordinating the field work and providing faecal samples. Thanks also to Jon Swenson and Gordon Luikart for comments on the manuscript.

References

Bellemain E, Swenson JE, Tallmon D, Brunberg S, Taberlet P (2004) Estimating population size of elusive animals with DNA from

- hunter-collected feces: comparing four methods for brown bears. *Conservation Biology* (in press).
- Miller SD, Joyce P, Waits LP (2002) Assessing allelic dropout and genotype reliability using maximum likelihood. *Genetics*, **160**, 357–366.
- Mills LS, Citta JJ, Lair K, Schwartz M, Tallmon D (2000) Estimating animal abundance using non-invasive DNA sampling: promise and pitfalls. *Ecological Applications*, **10**, 283–294.
- Paetkau D (2003) An empirical exploration of data quality in DNA-based population inventories. *Molecular Ecology*, **12**, 1375–1387.
- Paetkau D, Strobeck C (1994) Microsatellite analysis of genetic variation in black bear populations. *Molecular Ecology*, **3**, 489–495.
- Piggott M, Bellemain E, Taberlet P, Taylor A (2004) A multiplex pre-amplification method that significantly improves microsatellite amplification and error rates for faecal DNA in limiting conditions. *Conservation Genetics*, **5**, 417–420.
- Schwartz MC, Tallmon D, Luikart G (1999) Using genetics to estimate the size of wild populations: many methods, much potential, uncertain utility. *Animal Conservation*, **2**, 321–323.
- Smith KL, Alberts SC, Bayes MK *et al.* (2000) Cross-species amplification, non-invasive genotyping, and non-mendelian inheritance of human STRPs in savannah baboons. *American Journal of Primatology*, **51**, 219–227.
- Taberlet P, Luikart G (1999) Non-invasive genetic sampling and individual identification. *Biology Journal of the Linnean Society*, **68**, 41–55.
- Taberlet P, Griffin S, Goossens B *et al.* (1996) Reliable genotyping of samples with very low DNA quantities using PCR. *Nucleic Acids Research*, **24**, 3189–3194.
- Taberlet P, Camarra J-J, Griffin S *et al.* (1997) Noninvasive genetic tracking of the endangered Pyrenean brown bear population. *Molecular Ecology*, **6**, 869–876.
- Ulrich PP, Romeo JM, Daniel LJ, Vyas GN (1993) An improved method for the detection of hepatitis C virus RNA in plasma utilizing heminested primers and internal control RNA. *PCR Methods and Applications*, **2**, 241–249.
- Waits LP, Taberlet P, Swenson JE, Sandegren F (2000) Nuclear DNA microsatellite analysis of genetic diversity and gene flow in the Scandinavian brown bear (*Ursus arctos*). *Molecular Ecology*, **9**, 421–431.
- Waits LP, Luikart G, Taberlet P (2001) Estimating the probability of identity among genotypes in natural populations: cautions and guidelines. *Molecular Ecology*, **10**, 249–256.